

The association between Dairy Intake, Simple Sugars and Body Mass Index with Expression and Extent of Anger in Female Students

Naser Kalantari, MD¹

Saeid Doaei, Msc²

Maedeh Gordali, Bs³

Ghazal Rahimzadeh, Msc⁴

Maryam Gholamalizadeh, Msc⁵

1. Department of Nutrition and Food Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Student's Research Committee, National Nutrition and Food Technology Research Institute, faculty of Nutrition Sciences and Food Technology, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

3. Department of Community Nutrition, Shahid Beheshti University of Medical Sciences, Tehran, Iran

4. Department of Food Science and Technology, Science and Research Branch, Azad University, Tehran, Iran

5. Food Security Research Center, Department of Clinical Nutrition, School of Nutrition and Food Sciences, Isfahan University of Medical Sciences, Isfahan, Iran

Corresponding author:

Saeid Doaei,

Faculty of Nutrition Sciences and Food Technology, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Tel: +98 21 66902061

Fax: +98 21 66902038

Email: sdoaei@sbmu.ac.ir

Objective: A significant increase in violence in the world and its impact on public health and society can be an important reason to offer solutions to reduce or control anger. Studies have shown that specific food groups may be effective in controlling mental disorders such as depression, anxiety and anger. The purpose of this study was to determine the relationship between food intake and Body Mass Index on state-trait anger expression in female students of Shahid Beheshti University of Medical Sciences.

Method: In this cross-sectional study, 114 female students were randomly selected from dormitories of Shahid Beheshti University of Medical Sciences. Body height and weight were measured using the scale and stadiometer, respectively. The required data for evaluating the relationship between state-trait anger expression and food consumption groups were collected using State-Trait Anger Expression Inventory-2 (STAXI-2) and Food Frequency questionnaires.

Results: The results revealed a significant negative correlation between consumption of dairy product and trait anger (angry reaction), ($P = 0.015$). This association remained significant after adjustment of confounding factors. No significant correlations were found between other food groups as well as BMI and state-trait anger expression.

Conclusion: The higher intake of dairy products reduced state-trait anger expression. This result is consistent with the findings of many studies on the effect of dairy consumption on mental disorders. Therefore, consumption of dairy products can be a solution for reducing anger.

Key words: Anger, Body Mass Index, Food intake

Iran J Psychiatry 2016; 11:1: 43-50

Anger has been characterized as a complex construct including hostility, irritability and aggressive behavior (1). Anger can be defined as an emotional response or internal feeling which is caused by physiological excitation, cognition and malice thoughts (2). Nevertheless, anger is an emotion of satisfaction that prepares us to face the potential surrounding risks. On the other hand, destructive impact of anger may have detrimental effects on humans and their environments (3). Anger is a robust risk factor for Coronary Heart Disease (CHD) (4, 5), Cancer (6), and

Eating disorders (7) which can affect the quality of life (8, 9). Demark et al. (2011) have reported that over 36% of patients with brain injury have shown anger and aggression states (10). The effects of diet have been reported on depression (11), fervour (12), brain function (13, 14), stress response system (15), and oxidative reactions (16). On the other hand, many people eat to overcome emotional problems, mental pressure and anxiety (17, 18). There has been a positive correlation between obesity and some mental disorders including depression and anxiety (19, 20). Eating disorders in response to mental stress, fatigue,

anxiety, stress and depression have been reported (21, 22).

Many studies found that lack of micronutrients such as Zinc (23), vitamin E, Iron, Calcium, Iodine, Selenium, Folic acid (24), vitamin C (25), Choline (26), and vitamin D (27) is common among students all over the world. On the other hand, residency in dormitories can exacerbate the nutritional risk factors among students. Previous study showed that nutritional status of students living in the dorms were different from other students. Food consumption in dormitory students was decreased when entering university (28) and significant drop in food intake quality was observed in half of the dormitory students (29). Furthermore, despite the lack of some micronutrients intake, the nutritional status of students who were not living in the dorms were more acceptable compared to the dormitory students (30).

Previous studies about the effects of food and nutrient intake on mental disorders illustrated that a high intake of low-fat dairy products (31), decrease in consumption of fast food (32), using the Mediterranean Diet (33), and consumption of omega-3 fatty acids (34) and magnesium (35) supplements can result in reducing depression, anxiety and stress.

Further studies that investigated the relationship between food consumption and mental conditions have considered the prevalence of depression and its relation to food intake (36, 37). Although anxiety has been a common problem among students (38), no study was conducted on the relationship between diet and stressful situations (such as anger). Considering the results of studies conducted on the relationship between dietary intakes and mental disorders, we chose two food groups for evaluation.

Therefore, the purpose of this research was to determine the relation between dietary intake of dairy foods, simple sugars and BMI with states of anger in female students of a dormitory in Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Materials and Method

In this cross-sectional study, 114 resident students accommodating Fatima Zahra dormitory at Shahid Beheshti University of Medical Sciences were randomly selected and examined. This study was conducted from spring 2014 to autumn 2015.

Initially, students' weight and height were measured using Seca Scales (precision of 1 kg) and a tape measure (accuracy of 1 cm), respectively. After calculating the Body Mass Index, each participant was asked to complete the Anger Questionnaire, designed by Spielberger, validated by Barabadi in Iran, to measure anger in teens and adults (39). Spielberger and Brenner introduced this scale in 2003 and published it in 2009 to evaluate experience, expression, control and taming of anger. This questionnaire consists of 35 items, 4 subscales and 3 sections; the first part has 10 items, the second part contains 10

items, and the third part has 15 items, which evaluate anger state, anger trait and expression and control of anger, respectively. All sections of the questionnaire were calibrated using 3-point Likert scale.

To score the first section, the statements were identified as "never, a little, and very high" and scored as 1, 2, 3, respectively. To score the second section and 5 questions of the third section (which represents the expression of anger), the statements were identified as "very little, sometimes, and often" and scored as 1, 2, 3, respectively. To score the remaining questions in the third section (which represent the control of anger), scoring was done inversely, namely the score of "very little" was 3, the statement of "sometimes" was scored as 2, and the phrase "often" was awarded a score of 1 (40). The first part with the phrase "I feel right now ..." measured the state of anger (41) which graded the intensity of feeling. This section with 10 items included the anger state and subscales cases as follows: (a) Sense of Furiously (42), (b) Willingness to Verbal Expression of Anger (43), and (c) Trend to Physical Expressions of Anger (44). The second section, entitled "I feel usually...." Involved 10 items to measure trait anger (45), and was graded as the first part. The scale of trait anger has two subscales of angry state (46) and angry reaction (47). The title of the third section was "What is your behavioral reactions in angry mood?" that measured the expression and control of anger, and included four scales that involved subscales as follows: (a) Outer anger impart (48), (b) inner's anger impart (49), c: Control of outer anger (50) and (d) control of inner anger (51).

The food frequency questionnaire was used to investigate the intake of dairy food groups and simple sugars. This questionnaire was used to assess the frequency of food intake in a day, week or month. To facilitate the evaluation of this method, foods were organized as groups with similar nutrients. Since Food Frequency Questionnaire is based on frequency of food intake (not only specific nutrient), the obtained information was related to specific food group not specific nutrients (52). All participants signed the informed consent to participate in the study.

Linear regression was used for data analysis. Data were recorded as mean±standard deviation (SD). A statistical package (SPSS, version 21.0 for Windows, SPSS Inc.) was used for data processing. Differences were considered significant at $p < 0.05$.

Results

The personal characteristics of the study population are shown in Table 1. In the current study, 70% of the population aged between 20 to 25 years old. With respect to the academic degree, they were divided into three groups of Bachelor, MSc, and PhD. The number of students studying in each of these three groups was equal. In this community, the BMI was 21.61 ± 2.79 that was in the normal range. 13% of the Body Mass Index was less than 18.5; and 75% of the Body Mass

Index was between 18.5 to 24.9, and 12% of the body mass index was greater than or equal to 25, which was considered as overweight or obese.

With regards to anger measurements, for the first section, the mean of calculating the score was 13 out of 30. For trait anger, the calculated score was a little more than anger state (18.75 of 30). On the other hand, the resulting score for the third section (anger expression) was relatively high (28.35 of 45).

With respect to food groups intake, the average consumption of dairy products in this study was approximately 4 servings per day. According to the values recommended for adults, intake of dairy products should be at least 2 servings per day (50). Therefore, the dairy intake of the study population was averagely more than the minimum recommended values. The mean intake of simple sugars was approximately 7 servings per day that was higher than allowable values (0-3 servings) (50).

In this study, a significant inverse relationship was found between dairy consumption and trait anger (reacted furiously and angry state) ($P = 0.015$). On the other hand, there was not a significant correlation between dairy intake and simple sugars with other subscales of anger (state and expression of anger). Furthermore, there was a negative relevance between age and anger trait by investigating each parameters of age, height, weight, and Body Mass Index and its relationship with anger ($P=0.028$). Overall, age increase from 20 to 36 reduced levels of trait anger.

No significant relevance was found between Body Mass Index and other subscales of anger (state and expression of Anger). The relevance between dairy consumption, simple sugars, age and anthropometric indices with all anger subscales is demonstrated in Table 2.

One of the confounding variables in this study was supplements intake by students. To eliminate the effect of confounding variables and measuring the effect of calcium supplementation on the relationship between anger and dairy intake, all persons who were consuming calcium supplements or multivitamin-mineral (14) were excluded from the study, and the results were reviewed. After the removal of the confounding variables, the relationship between dairy consumption and trait anger remained significant ($P = 0.027$). Table 3 displays the relationship between dairy consumption and trait anger after removing the confounding variables.

Discussion

There was a negative correlation between dairy intake (on levels recommended by the Food Guide Pyramid) and trait anger (reacted furiously and angry state) (P -value = 0.015). Therefore, students who had more dairy products were less in angry states and had low angry reactions. Miyake et al. reported similar results and showed that more consumption of yogurt and calcium

was associated with less rampancy of depression symptoms (53). Crichton et al. reported that consumption of some low-fat dairy products had beneficial effects on social functioning, stress and memory (54). Moreover, Roohafza et al. found that more consumption of dairy products respond to low stress levels (55). This effect may be due to the high content of tryptophan (essential amino acid) in dairy products. Tryptophan is a precursor for serotonin (a neurotransmitter in brain) that has a significant effect on mental states. In most previous studies, levels of brain's serotonin in patients with depression and anxiety were lower than healthy subjects (56). Duke (57) and Giessen et al. (58) showed that serotonin function has an inverse correlation with the degree of anger and hostility.

Moreover, this effect can be related to the antioxidant content of dairy products such as vitamins A and B2¹ (57). Antioxidants can affect oxidative stress, which may lead to slower neuronal changes that are associated with depression in old age (59).

In this research, no significant correlation was found between intake of simple sugars, and the expression and extend anger subscales; these results were also confirmed in other studies. On the other hand, Keith et al. found that low-carbohydrate diet in female group is associated with depression and anger (60). In the current study, consumption of simple sugars (not total carbohydrate) in the long-term (1 year) was evaluated, and these factors may cause the differences in the results. There was not a significant effect of Body Mass Index on expression and extent of observed anger.

Similarly, Arian et al. (61) and Ahmadi et al. (62) investigated that the prevalence of mental health problems had no significant relationship with anthropometric indices (including BMI).

No association was found between BMI and depression in any other similar studies (63-65). However, in some cases, a significant positive correlation was detected between depression and Body Mass Index (66-69). These differences in the results can be related to several factors such as age, gender, socioeconomic and cultural factors that affect this relationship. Furthermore, social attitudes to obesity should be considered as a important cultural factor (70). For example, in Western culture, negative attitudes toward obesity can influence self-esteem and cause depression (71). While in some other societies larger body size is accepted (72).

¹ Vitamin B2 is one of many nutrients required to recycle glutathione, which is one of the most important antioxidants in the human body. From a chemical standpoint, what B2 does is facilitate the conversion of oxidized glutathione into reduced glutathione.

Limitations

Since only female students and limited dormitories were evaluated in this study, conducting similar research with larger samples is required. In addition, investigation of other food groups associated with anger is a suggestion that can be offered in relation to this research. The findings of this study are based on cross-sectional data, and an important avenue for future research will be to replicate these studies with causal structures using longitudinal data.

Conclusion

Life quality can be affected by mental health as well as physical health. According to the results of this study, more consumption of dairy food group decreases the trait anger. Since the subscale of trait anger included angry state and angry reaction, it can be concluded that dairy products intake has a positive effect on both temperament during long time and immediate reaction to oppositions. Therefore, intake of dairy products can be suggested as a nutritional strategy for reducing anger and increasing soothe.

Acknowledgement

We thank all of the participants and questionnaire respondents for their contribution to this study. This work was supported by the Student's Research Committee of Shahid Beheshti University of Medical Sciences, Iran (grant number 5632).

Conflict of interest

There is no conflict of interest to be declared.

Table 1: The Basic Characteristics of the Participants

Index	Mean	Standard deviation
High (m)	1.63	0.047
Weigh (kg)	58.12	7.95
Body mass index (kg/m ²)	21.61	2.79
Anger state (score of 30)	12.98	3.53
Trait Anger (score of 30)	18.75	4.01
Anger expression (score of 45)	28.37	6.36
Dairy consumption (serving per day)	3.89	2.06
Intake sugar (serving per day)	7.36	6.52

Table 2: Relationship between Dairy Consumption and Simple Sugars, Age and Anthropometric Indices with Anger

Index	Anger State			Trait anger			Anger expression		
	B Coefficient	Beta coefficient	P-value	B Coefficient	Beta coefficient	P-value	B Coefficient	Beta coefficient	P-value
Dairy intake	0.000	0.032	0.753	-0.001	-0.23	0.015*	-0.001	-0.105	0.273
Sugar intake	-6.42	-0.043	0.651	0.000	0.062	0.5	-8.80	-0.033	0.729
Age	-0.84	-0.074	0.445	-0.271	-0.210	0.028*	0.091	0.044	0.640
Body mass index	1.636	1.292	0.472	-0.391	-0.272	0.877	-7.021	-3.075	0.083
High	42.23	0.561	0.485	-12.05	-0.141	0.858	-190.004	-1.401	0.079
Weight	-0.594	-1.333	0.481	0.176	0.348	0.851	2.521	3.140	0.094

Table 3: The Relationship between Dairy Consumption with Trait Anger after Excluding 14 Patients

Index	B Coefficient	Beta coefficient	P-value
Dairy intake	-0.001	-0.221	0.027*

References:

1. Mick E, McGough J, Deutsch CK, Frazier JA, Kennedy D, Goldberg RJ. Genome-wide association study of proneness to anger. *PLoS one* 2014; 9: e87257.
2. O'Neill H, *Managing anger*, New York: Wiley; 2006.
3. Taylor JL, Novaco RW. *Anger treatment for people with developmental disabilities: A theory, evidence and manual based approach*. City: John Wiley & Sons; 2005.
4. Masoudnia E. Hostility, anger and risk of coronary artery atherosclerosis. *SSU_Journals* 2011; 18: 540-551.
5. Bleil ME, McCaffery JM, Muldoon MF, Sutton-Tyrrell K, Manuck SB. Anger-related personality traits and carotid artery atherosclerosis in untreated hypertensive men. *Psychosomatic medicine* 2004; 66: 633-639.
6. Geyer S. Life events prior to manifestation of breast cancer: a limited prospective study covering eight years before diagnosis. *J Psychosom Res* 1991; 35: 355-363.
7. Miotto P, Pollini B, Restaneo A, Favaretto AA, Preti A. Aggressiveness, anger and hostility in eating disorders. *Comprehensive psychiatry* 2008; 49: 364-373.
8. Ahmadi F, GhofraniPour F, Arefi H, Abedi H, Faghihzade S. The effect of counselling on quality of life of coronary heart disease patients. *J Psychol* 2000; 4: 97-102.
9. Li R, Yan BP, Dong M, Zhang Q, Yip GW-K, Chan C-P, et al. Quality of life after percutaneous coronary intervention in the elderly with acute coronary syndrome. *International journal of cardiology* 2012; 155: 90-96.
10. Demark J, Gemeinhardt M. Anger and its management for survivors of acquired brain injury. *Brain Injury* 2002; 16: 91-108.
11. Jacka FN, Pasco JA, Mykletun A, Williams LJ, Hodge AM, O'Reilly SL, et al. Association of Western and traditional diets with depression and anxiety in women. *American Journal of Psychiatry* 2010.
12. Liu S, Manson JE, Buring JE, Stampfer MJ, Willett WC, Ridker PM. Relation between a diet with a high glycemic load and plasma concentrations of high-sensitivity C-reactive protein in middle-aged women. *The American journal of clinical nutrition* 2002; 75: 492-498.
13. Gómez-Pinilla F. Brain foods: the effects of nutrients on brain function. *Nature Reviews Neuroscience* 2008; 9: 568-578.
14. Molteni R, Barnard R, Ying Z, Roberts C, Gomez-Pinilla F. A high-fat, refined sugar diet reduces hippocampal brain-derived neurotrophic factor, neuronal plasticity, and learning. *Neuroscience* 2002; 112: 803-814.
15. Tannenbaum BM, Brindley DN, Tannenbaum GS, Dallman MF, McArthur MD, Meaney MJ. High-fat feeding alters both basal and stress-induced hypothalamic-pituitary-adrenal activity in the rat. *American Journal of Physiology-Endocrinology and Metabolism* 1997; 273: E1168-E1177.
16. Engelhart MJ, Geerlings MI, Ruitenberg A, van Swieten JC, Hofman A, Witteman JC, et al. Dietary intake of antioxidants and risk of Alzheimer disease. *Jama* 2002; 287: 3223-3229.
17. Macht M. How emotions affect eating: a five-way model. *Appetite* 2008; 50: 1-11.
18. Nishitani N, Sakakibara H, Akiyama I. Eating behavior related to obesity and job stress in male Japanese workers. *Nutrition* 2009; 25: 45-50.
19. Simon GE, Von Korff M, Saunders K, Miglioretti DL, Crane PK, van Belle G, et al. Association between obesity and psychiatric disorders in the US adult population. *Archives of general psychiatry* 2006; 63: 824-830.
20. Ahmadi SM, Keshavarzi S, Mostafavi S-A, Lankarani KB. Depression and Obesity/Overweight Association in Elderly Women: a Community-Based Case-Control Study. *Acta Medica Iranica* 2015; 53.
21. Kandiah J, Yake M, Jones J, Meyer M. Stress influences appetite and comfort food preferences in college women. *Nutrition Research* 2006; 26: 118-123.
22. Ranjbar E, Kasaei MS, Shirazi MM, Nasrollahzadeh J, Rashidkhani B, Shams J. Effects of Zinc Supplementation in Patients with Major Depression: A Randomized Clinical Trial. *Iran J Psychiatry*. 2013; 8: 73-79.
23. Oliveras LM, Nieto GP, Agudo AE, Martínez MF, López GdISH, López MM. [Nutritional assessment of a university population]. *Nutricion hospitalaria* 2005; 21: 179-183.
24. Gan R, Eintracht S, Hoffer LJ. Vitamin C deficiency in a university teaching hospital. *Journal of the American College of Nutrition* 2008; 27: 428-433.
25. Gossell-Williams M, Benjamin J. Choline: are our university students eating enough? *West Indian medical journal* 2006; 55: 197-199.
26. Nakamura K, Nashimoto M, Tsuchiya Y, Obata A, Miyanishi K, Yamamoto M. Vitamin D insufficiency in Japanese female college students: a preliminary report. *International journal for vitamin and nutrition research* 2001; 71: 302-305.
27. Rezaei-pour A, Yavari P, Mahmodi M, Fili S. The function of female medical student in prevention of iron deficiency anemia due to the menarche. *Hayat* 2003; 8: 50.
28. Amani R. Assessment of nutrition related life style patterns of female students in Ahvaz universities dormitories. 2004; 42: 54-61.
29. esfanjani AT, Mahdavi R, Ghaemmaghami J, Safaeian A. Comparison between students who resident in dormitory and who are not in Ardabil university medical of science. *Ardabil University of Medical Science Journal* 2003; 9: 44.
30. Crichton GE, Murphy KJ, Bryan J. Dairy intake and cognitive health in middle-aged South Australians. *Asia Pacific journal of clinical nutrition* 2010; 19: 161-171.

31. Bakhtiyari M, Ehrampoush E, Enayati N, Rastmanesh R, Delpisheh A, Zayeri F, et al. Correlation Between Fast Food Consumption And Levels Of Anxiety In Students Of Medical Science Universities In Tehran. 2011.
32. Oliveira C, Oliveira G, Espildora F, Girón R-M, Vendrell M, Dorado A and et al. Mediterranean diet is associated on symptoms of depression and anxiety in patients with bronchiectasis. *General hospital psychiatry* 2014; 36: 277-283.
33. Stoll AL, Severus WE, Freeman MP, Rueter S, Zboyan HA, Diamond E, et al. Omega 3 fatty acids in bipolar disorder: a preliminary double-blind, placebo-controlled trial. *Archives of general psychiatry* 1999; 56: 407-412.
34. Abbasi B, Kimiagar M, Shirazi MM, Sadeghnia kh, Rashidkhani B, Karimi N, et al. Effect of Magnesium Supplementation on Physical Activity of Overweight or Obese Insomniac Elderly Subjects: A Double-Blind Randomized Clinical Trial. *Zahedan Journal of Research in Medical Sciences* 2013; 15: 14-19.
35. Appleton KM, Peters TJ, Hayward RC, Heatherley SV, McNaughton SA, Rogers PJ, et al. depressed mood and n-3 polyunsaturated fatty acid intake from fish: non-linear or confounded association? *Social psychiatry and psychiatric epidemiology* 2007; 42: 100-104.
36. Timonen M, Horrobin D, Jokelainen J, Laitinen J, Herva A, Räsänen P. Fish consumption and depression: the Northern Finland 1966 birth cohort study. *Journal of affective disorders* 2004; 82: 447-452.
37. Eisenberg D, Gollust SE, Golberstein E, Hefner JL. Prevalence and correlates of depression, anxiety, and suicidality among university students. *American Journal of Orthopsychiatry* 2007; 77: 534-542.
38. Barabadi A, Heydarinasab L. Preliminary validation of the questionnaire State-Trait Anger Expression Inventory -2 Child and Adolescent (STAXI-2 C/A) amongst a sample of Iranian students aged 12-17 years. *Journal of Sabzevar University of Medical Sciences* 2013; 20: 281-291.
39. Spielberger CD. State-Trait Anger Expression Inventory (S.T.A.X.I.) Professional Manual. Psychological Assessment Resources, New York: Wiley; 1988.
40. Redford W. Anger kills. Translated by Kadkhodaeian M and ZarrinKalak SH. Tehran, Firooze, 2000.
41. Mayer JD, Caruso DR, Salovey P. Emotional intelligence meets traditional standards for an intelligence. *Intelligence* 1999; 27: 267-298.
42. Brackett MA, Mayer JD, Warner RM. Emotional intelligence and its relation to everyday behaviour. *Personality and Individual differences* 2004; 36: 1387-1402.
43. Ramezani V, Abdollahi MH. The Study of the relationship between emotional intelligence and anger's incidence and controlling it among students. *Journal of psychology* 2006; 5: 66-83.
44. Julkunen J, Ahlström R. Hostility, anger, and sense of coherence as predictors of health-related quality of life. Results of an ASCOT substudy. *Journal of Psychosomatic Research* 2006; 61: 33-39.
45. Alan K. Positive psychology. Translated by Pasha Sharifi H and Najafizand J. Tehran: Sokhan; 2006.
46. Dahlen E, Deffenbacher J. Anger management: Empirically supported cognitive therapies; Current status and future promise. New York 2001.
47. Marshall JR. Motivation and Emotion, Translated by Seyyed Mohammadi Y. Tehran: Virayesh; 2006. 48-Smith BA, Kinney TA, Donzella B. The Influences of Gender, Self-Discrepancies and Self-Awareness on Anger and verbal Aggressiveness among US college students. *The Journal of Social Psychology*. 2001; 141: 245-275.
48. Agneta G, Fischer. R. Gender and emotion, social psychological perspective. London: Cambridge University Press; 2000.
49. Friedman M, Rosenman RH. Type a behavior and your heart. City: Knopf New York; 1974.
50. Mahan LK, Escott-Stump S. Krause's food & nutrition therapy, Philadelphia: Saunders/Elsevier; 2014.
51. Miyake Y, Tanaka K, Okubo H, Sasaki S , Arakawa M. Intake of dairy products and calcium and prevalence of depressive symptoms during pregnancy in Japan: a cross-sectional study. *BJOG: An International Journal of Obstetrics & Gynaecology* 2015; 122: 336-343.
52. Crichton GE, Murphy KJ, Bryan J. Dairy intake and cognitive health in middle-aged South Australians. *Asia Pacific journal of clinical nutrition* 2010; 19: 161.
53. Roohafza H, Sarrafzadegan N, Sadeghi M, Rafieian-Kopaei M. The association between stress levels and food consumption among Iranian population. *Archives of Iranian medicine* 2013; 16: 145.
54. Åsberg M, Träskman L , Thorén P. 5-HIAA in the cerebrospinal fluid: a biochemical suicide predictor? *Archives of general psychiatry* 1976; 33: 1193-1197.
55. Duke AA, Bègue L, Bell R, Eisenlohr-Moul T. Revisiting the serotonin-aggression relation in humans: A meta-analysis. *Psychological bulletin* 2013; 139: 1148.
56. Van de Giessen E, Rosell DR, Thompson JL, Xu X, Girgis RR, Ehrlich Y, et al. Serotonin transporter availability in impulsive aggressive personality disordered patients. *J Psychiatry Res* 2014; 58: 147-54.
57. Rahman A, Baker PS, Allman R , Zamrini E. Dietary factors and cognitive impairment in community-dwelling elderly. *The journal of nutrition, health & aging* 2007; 11: 49.
58. Joseph JA, Shukitt-Hale B, Denisova NA, Bielinski D, Martin A, McEwen JJ, et al. Reversals of age-related declines in neuronal signal transduction, cognitive, and motor behavioral deficits with blueberry, spinach, or strawberry dietary supplementation. *The Journal of Neuroscience* 1999; 19: 8114-8121.

59. Keith RE, O'Keeffe KA, Blessing DL, Wilson GD. Alterations in dietary carbohydrate, protein, and fat intake and mood state in trained female cyclists. *Medicine & Science in Sports & Exercise* 1991.
60. Arian V, Eftekhari M.H, Tabatabaei H.R, Naziri Q. The study of the prevalence of depression in male teachers 25-50 years old in the city of Shiraz and it's relation with anthropometric indicators at year 2006, comprehensive portal of congresses. University of Medical Sciences of Tabriz, 2006.
61. Ahmadi S.M, Eftekhari M.H, Firoozabadi A, RajaeiFard A.R, Sawid M. The study of severity and type of obesity in depressed fat women (15-49) years old referring to Shahid Motahhari clinic in Shiraz, comprehensive portal of congresses, university of Medical Sciences of Tabriz , 2006.
62. Roohafza H, Kelishadi R, Sadeghi M, Hashemipour M, Pourmoghaddas A , Khani A. Are obese adolescents more depressed? *Journal of education and health promotion* 2014; 3.
63. Mikkilä V, Lahti-Koski M, Pietinen P, Virtanen SM , Rimpelä M. Associates of obesity and weight dissatisfaction among Finnish adolescents. *Public health nutrition* 2003; 6: 49-56.
64. Wadden TA, Foster GD, Stunkard AJ, Linowitz JR. Dissatisfaction with weight and figure in obese girls: discontent but not depression. *International journal of obesity* 1988; 13: 89-97.
65. Raf'ati F, Pilevarzade M, Mohammadi Soleimani M.R, Salari S, PormehrYabande A. The association of anthropometric indicators with depression and mental health at nursing students in Jiroft. *Journal of endocrinology and metabolism of Iran* 2012; 1: 135-141.
66. Geoffroy M-C, Li L, Power C. Depressive symptoms and body mass index: co-morbidity and direction of association in a British birth cohort followed over 50 years. *Psychological medicine* 2014; 44: 2641-2652.
67. Wang H, Fu J, Lu Q, Tao F, Hao J. Physical activity, body mass index and mental health in Chinese adolescents: a population based study. *The Journal of sports medicine and physical fitness* 2014; 54: 518-525.
68. Sörberg A, Gunnell D, Falkstedt D, Allebeck P, Åberg M , Hemmingsson T. Body mass index in young adulthood and suicidal behavior up to age 59 in a cohort of Swedish men. *PloS one* 2014; 9: e101213.
69. Wardle J, Cooke L. The impact of obesity on psychological well-being. *Best Practice & Research Clinical Endocrinology & Metabolism* 2005; 19: 421-440.
70. Williams K, Taylor CA, Wolf KN, Lawson R , Crespo R. Cultural perceptions of healthy weight in rural Appalachian youth. *Rural Remote Health* 2008; 8: 932.
71. Bergstrom RL, Neighbors C, Lewis MA. Do men find "bony" women attractive? Consequences of misperceiving opposite sex perceptions of attractive body image. *Body Image* 2004; 1: 183-191.
72. Wardle J, Cooke L. The impact of obesity on psychological well-being. *Best Practice & Research Clinical Endocrinology & Metabolism* 2005; 19: 421-440.