

The Prevalence of Postpartum Depression and Its Association with Food Insecurity among Mothers Referring to Community Health Centers

Neda Ezzeddin^{1*}, Hassan Jahanihashemi², Roza Zavoshy³, Mostafa Noroozi²

Abstract

Objective: Postpartum depression (PPD) is a condition which may compromise both maternal and neonatal health. The present study was conducted to determine the prevalence of PPD and its association with demographic, socioeconomic, obstetric, and household food security status.

Method: This cross sectional study was conducted in community health centers in west of Tehran. A total of 325 women were selected by stratified sampling method from community health centers. A sociodemographic questionnaire, USDA 18-item questionnaire, and Edinburgh Postnatal Depression Scale (EPDS) questionnaire were used for data collection. Data were analyzed using both descriptive and analytic analyses, such as chi-squared test and logistic regression in SPSS 22.

Results: The prevalence of PPD and food insecurity among the studied population was 35.4% and 34.2%, respectively. The results of this study revealed a significant association among PPD and food insecurity (OR = 6.690, CI = 3.118-14.353, $p < 0.001$), the levels of economic satisfaction (OR = 3.419, CI = 1.241-9.420, $P = 0.017$), pregnancy loss (OR = 1.899, CI = 1.006-3.582, $p = 0.048$), and pregnancy complications (OR = 1.853, CI = 1.083-3.170, $P = 0.024$).

Conclusion: Based on the results of this study, household food insecurity may predispose mothers to PPD. Moreover, it was observed that mothers with poor economic satisfaction were more likely to be depressed. Histories of pregnancy loss and pregnancy complications were other risk factors for PPD.

Key words: Food Security, Pregnancy Complication, Postpartum Depression, Socioeconomic

Postpartum depression (PPD), whose onset is generally within 12 weeks after delivery, is a disorder that is often unrecognized and undertreated (1, 2). PPD is characterized by sadness, fatigue, irritability, and disinterest in life events (3). PPD may compromise both maternal and neonatal health in different ways, such as infant development, breast feeding, sleep routines, vaccinations, and safety practice (4, 5). Economic, social and biological factors, obstetrical history, lifestyle and history of mental illness are identified as risk factors for PPD (6).

Unwanted pregnancy, due to lack of family planning, may increase the prevalence of PPD (7). Giannandrea et al. (8) found that women with the history of loss of pregnancy (miscarriage, stillbirth, or induced abortion) were more likely to be diagnosed with psychiatric disorder, particularly depression.

Social factors, such as income inequality increases the risk of PPD (9). Studies showed that poor women experience more food insecurity compared to higher income groups (10). Household food security is defined as accessibility of enough, safe, and good-quality food for all individuals at all times to meet their dietary needs

1. Department of Community Nutrition, School of Nutrition Science and Food Technology, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

2. Children Growth Research Center, Qazvin University of Medical Sciences, Qazvin, Iran.

3. Department of Nutrition, Children Growth Research Center, Qazvin University of Medical Sciences, Qazvin, Iran.

***Corresponding Author:**

Address: No. 7, Shahid Farahzadi Blvd, Shahid Hafezi St. (Western Arghavan), Ghods Town (West), Tehran, Iran.
Tel: 98-2122077424, Fax: 98-2122360660, Email: Neda.Ezzeddin@sbm.ac.ir

Article Information:

Received Date: 2017/03/18, Revised Date: 2018/5/25, Accepted Date: 2018/07/01

to support a healthy and productive life (11); and food insecurity refers to an opposite condition and is related to poverty and low-income (12). Food insecurity is an important problem in the developing world and has impacts on nutrition status, growth, and development (13). The association of food insecurity with maternal depression and decreased mental health status has been reported in several studies (12, 14–16). However, a few studies have assessed the association between PPD and food insecurity (17). In Iran, most studies have examined the association between PPD and mothers' diet (18, 19) or the association between food security and maternal depression (not PPD) (20). Therefore, the present study was conducted to determine the prevalence of PPD and its related risk factors, with emphasis on food insecurity, in a small population of Iranian mothers.

Materials and Methods

1. Study Population, Design, and Data Collection

This cross sectional study was performed from March to June 2014 in community health centers in the west of Tehran. The population of the study included postpartum women (3 to 8 months postdelivery), aged 18–45 years, with no history of depression (self-reported and according to didn't visit a psychologist or psychiatrist for depression) and chronic disease (such as cardiovascular disease, diabetes, cancers and endocrine disease). In this study, all women agreed to participate in the study and signed the informed consent form. (The ethical code: D.44.21039).

A total of 8 community health centers in the west of Tehran, which provided a good distribution of all 16 centers based on covered population and geographic area, were randomly selected. Then, using stratified sampling technique (proportional to size), 325 women were randomly selected from these community health centers.

2. Measures

2-1. Food Security Assessment

Food security status was evaluated using the USDA (US Department of Agriculture) questionnaire (21). This 18-item questionnaire, which assessed household food security status, was completed by interviewing women. Scoring the USDA 18-item questionnaire is as follows: responses "most of the time correct" and "sometimes correct" in questions 1 to 3 and 11 to 13, "almost every month" and "some months" in questions 5, 10, and 16, and "yes" in questions 4 and 6 to 9, 14, 15, 17, and 18 are scored 1; and responses "is not correct", "does not know or avoids", "only once or twice a month", and "no" are scored 0. The final score is calculated based on the number of positive responses.

The studied women were divided into 4 groups based on the acquired scores of the questionnaire: (1) food secure, (2) food insecure without hunger, (3) food insecure with moderate hunger, and (4) food insecure with severe hunger (Table 1) (22). The 2 groups of "food insecure with moderate hunger" and "food insecure with severe

hunger" were merged together and formed the food insecure with hunger group. The validity of this questionnaire has been evaluated in another study on Iranian households in Shiraz and results demonstrated acceptance of the questionnaire (23).

2-2. Postpartum Depression Assessment

PPD was assessed by the Edinburgh Postnatal Depression Scale (EPDS) questionnaire. EPDS is a 10-item self-report scale for measuring the intensity of depressive symptoms experienced within the past 7 days, and each item is rated on a 4-point scale from 0 to 3. The cut-off point for PPD is 12/13 or greater. Scores of 10 to 12 represent borderline and 0 to 9 non-depressed women (24). In this study, we used the Persian version of EPDS with cut-off of 13 and reliability of 0.83 (Cronbach's alpha) (25).

2-3. Assessment of Demographic, Socio economic, and Obstetric Status

Demographic and socio-economic characteristics (including maternal and paternal age; education; occupational status; and level of economic satisfaction) and obstetric information (including pregnancy ranks, history of pregnancy loss: miscarriage, stillbirth, or induced abortion; unwanted infant gender; unwanted pregnancy; type of delivery; and pregnancy complications, such as gestational diabetes mellitus, pre-eclampsia, anemia, high blood pressure, nausea and vomiting, premature contractions, and bleeding) were collected via the questionnaire.

To assess the levels of economic satisfaction, mothers were asked to choose a number from 1 to 10 (subjectively) to describe their satisfaction. For data analysis, the scores were placed in 3 categories: (a) low satisfaction, 1 to 4, (b) moderate satisfaction, 5 to 7, and (c) high satisfaction, 8 to 10.

3. Statistical Analysis

The association between PPD and qualitative (pregnancy ranks, type of delivery, history of pregnancy loss, unwanted infant gender, unwanted pregnancy, parents occupational status, parents educational levels, levels of economic satisfaction, pregnancy complications, and household food security status) and quantitative (maternal age) variables were evaluated with chi-squared and t tests, respectively, in SPSS22. Also, the logistic regression was used to control the variables that were statistically associated with PPD. The significance level was set at 0.05.

Results

The mean age of participants was 28.62 ± 5.67 years. Most of the mothers were housekeepers (86.5%), graduated (78.2%), and in the moderate level of economic satisfaction (64.1%). The prevalence of food insecurity and PPD among the studied women was 34.2% ($n = 111$) and 35.4% ($n = 115$), respectively. unwanted pregnancy (20.6%) and unwanted infant gender (13.7%) have also been reported, and the

prevalence of vaginal delivery was very low (22.2%). More detailed information about demographic, socioeconomic, and obstetric information is presented in Table 3.

The prevalence of food insecurity was significantly higher among depressed women and it was positively associated with food insecurity scores ($P < 0.001$) (Table). Among the examined variables, 8 variables (maternal education ($P = 0.013$), paternal education ($P = 0.010$), father employment ($P = 0.018$), economic satisfaction ($P < 0.001$), history of pregnancy loss ($P = 0.048$), unwanted pregnancy ($P < 0.001$), unwanted infant gender ($P = 0.004$) and pregnancy complications ($P < 0.001$) were significantly associated with PPD. However, there was no significant association among other variables (maternal age ($P = 0.404$), paternal age ($P = 0.121$), maternal occupational status ($P = 0.226$), type of delivery ($P = 0.070$), and pregnancy ranks ($P = 0.495$)) and PPD (Table 2).

Based on the results of logistic regression analyses, food insecurity, economic satisfaction, history of pregnancy loss, and pregnancy complications were significant ($P < 0.05$) (Table 4).

Discussion

The result of the study revealed a significant association among household food security status, levels of economic satisfaction, history of pregnancy loss, and pregnancy complications with PPD, but it did not have any significant association with other variables, including parents' age, parents' education level, parents' occupational status, pregnancy ranks, type of delivery (NVD or caesarean), unwanted pregnancy, and unwanted infant gender.

The prevalence of PPD in developed (high-income) countries is almost 13% and the prevalence of PPD may be even greater in developing countries (7). In this study, the prevalence of PPD among mothers was 35.4%. Khooshemehry conducted a study on mothers who referred to community health centers in the north of Tehran and found PPD prevalence of 30% (26). However, in the Manshoori's study, it was reported to be 68.5% among mothers who referred to community health centers in Rafsanjan (27), which is higher than our studied mothers. In another study by Abdollahi, the prevalence of PPD was 10.7% among mothers who referred to rural and urban primary health care centers of Mazandaran province (28). According to a meta-analysis and systematic review, the prevalence of PPD in Iran is 24.3%, based on the Edinburgh Postnatal Depression Scale; also, a significant geographic difference was observed in the prevalence of PPD (29).

The prevalence of pregnancy complications is higher among women in developing countries than in developed countries (30). In a cohort study on 4941 pregnant women, it was shown that pregnancy complications increased the risk of PPD (31). In this

study also, such an association was observed that was consistent with Shivalli and Blom studies (31, 32).

In the final logistic regression analyses, no significant association was found between unwanted pregnancy and PPD. This finding was consistent with the result of the study by Blom (31) and Sooki (33). However, in Barton (34) and Ria (35) studies, unwanted pregnancy was an important risk factor for PPD because it impose anticipated strain on families in meeting their basic needs (36).

The results of the present study revealed that household food insecurity is an important risk factor for PPD symptoms among mothers, which is consistent with the results of the study conducted by Dewing (17) and Patel (37). Food security is important for mothers' health, especially during pregnancy, because nutritional demands increase in this period (12). Studies have also indicated that food insecurity leads to maternal stress during pregnancy, and food-insecure pregnant mothers worry about having enough food for themselves and their family (38). Stressful life events put women at risk for PPD (39). According to some studies, depression could be alleviated by decreasing household food insecurity (15, 40). We just assessed food security among mothers, but we did not have any data on nutritional intakes and dietary biomarkers from blood samples or the levels of mother stress during pregnancy. Thus, we could not provide the mentioned reasons in other studies for our results, and they are just possible reasons for the observed associations.

Table 1. Classification of the Household Food Security Status Based on Scores

Household Food Security Status	Number of Positive Responses	
	Households without children under 18 years	Households with children under 18 years
Food secure	0-2	0-2
Food insecurity without hunger	3-5	3-7
Food insecurity with moderate hunger	6-8	8-12
Food insecurity with sever hunger	9-10	13-18

Table 2. The Association between Postpartum Depression and Quantitative Variables among Studied Mothers

Variables	Having PPD Mean±SD [*]			P-value**
	Yes	No	Total	
Mother age	28.97±6.37	28.42±5.25	28.62 ±5.67	0.404
Father age	33.89±6.73	32.80±5.60	33.19±6.04	0.121

^{*}SD= standard deviation

^{**}T test

Table 3. The Association between Postpartum Depression and Qualitative Variables among Studied Mothers

Variables	Having PPD N (%)			P-value*
	Yes	No	Total	
pregnancy ranks				
1	51(44.3)	106(50.5)	157(48.3)	0.495
2	38(33)	66(31.4)	104(32)	
>2	26(22.6)	38(18.1)	64(19.7)	
Total	115(100)	210(100)	325(100)	
type of delivery				
NVD ^{**}	19 (16.5)	53(25.2)	72(22.2)	0.07
Caesarean	96(83.5)	157(74.8)	253(77.8)	
Total	115(100)	210(100)	325(100)	
history of pregnancy loss				
Yes	31(27)	37(17.6)	68(20.9)	0.048
No	84(73)	173(82.4)	257(79.1)	
Total	115(100)	210(100)	325(100)	
Unwanted infant gender				
Yes	24(21.2)	20(9.6)	44(13.7)	0.004
No	89(78.8)	189(90.4)	278(86.3)	
Total	113(100)	209(100)	322(100)	
unwanted pregnancy				
Yes	38(33)	29(13.8)	67(20.6)	<0.001
No	77(67)	181(86.2)	258(79.4)	
Total	115(100)	210(100)	325(100)	
Mother occupational status				
Employment	12(10.4)	32(15.2)	44(13.5)	0.226
Housekeepers	103(89.6)	178(84.8)	281(86.5)	
Total	115(100)	210(100)	325(100)	

Father occupational Status				
Self-employed	87(76.3)	132(63.5)	219(68)	0.018
Government employee	27(23.7)	76(36.5)	103(32)	
Total	114(100)	208(100)	322(100)	
Education level of mother				
Under diploma	34(29.6)	37(17.6)	71(21.8)	0.013
diploma or higher	81(70.4)	173(82.4)	254(78.2)	
Total	115(100)	210(100)	325(100)	
Education level of father				
Under graduated	39(33.9)	44(21)	83(25.5)	0.010
graduate	76(66.1)	166(79)	242(74.5)	
Total	115(100)	210 (100)	325(100)	
Levels of economic satisfaction				
Poor	42(36.5)	32(15.4)	74(22.9)	<0.001
Moderate	65(56.5)	142(68.3)	207(64.1)	
Good	8(7)	34(16.3)	42(13)	
Total	115(100)	208(100)	323(100)	
Household food security status				
secure	51(44.3)	163(77.6)	214(65.8)	<0.001
Unsecure without hunger	24(20.9)	32(15.2)	56(17.2)	
Unsecure with moderate and severe hunger	40(34.8)	15(7.1)	55(16.9)	
Total	115(100)	210(100)	325(100)	
obstetric complications				
Yes	62(53.9)	68(32.4)	130(40)	<0.001
No	53(46.1)	142(67.6)	195(60)	
Total	115(100)	210(100)	325(100)	

* Chi-square test
 ** Normal Vaginal Delivery

Table 4. Final Logistic Regression Model for Predicting Postpartum Depression

variables	P-value	OR*	95 % C.I.**	
			Lower	Upper
Household food security status				
Food secure		Reference		
Food insecure without hunger	0.070	1.883	0.948	3.739
Food insecure with moderate and severe hunger	<0.001	6.690	3.118	14.353
Unwanted pregnancy	0.133	1.677	0.854	3.294
Paternal education	0.466	1.341	0.610	2.945
Maternal education	0.923	1.042	0.452	2.401
Economic satisfaction				
Poor		Reference		
Moderate	0.017	3.419	1.241	9.420

Good	0.153	1.926	0.784	4.733
Unwanted infant gender	0.360	1.470	0.644	3.356
Father occupational Status	0.213	1.486	0.797	2.770
pregnancy loss	0.048	1.899	1.006	3.582
obstetric complication	0.024	1.853	1.083	3.170

Note: table is based on results of binary logistic regression (only for significant situations

Based on chi-square test on table 3).

* Odds ratio **Confidence Interval

Limitation

The sample size in the study had enough statistical power to identify a significant association of PPD with household food security status, levels of economic satisfaction, and history of pregnancy loss and pregnancy complication; however, for other variables, a larger sample size may be required. In this study, we only assessed food security among mothers and we did not assess other nutritional considerations. Therefore, future studies with larger sample size and more nutritional assessments or even longitudinal studies may provide more accurate results. In this study, to better assess household food security, we assessed PPD risk factors in women with no history of referring to clinics for depression. Thus, the prevalence of PPD might have been underestimated. Another limitation was that this study was conducted in a small geographic area, so we could not generalize the results to all areas.

Conclusion

Household food insecurity may predispose mothers to PPD. We also observed that mothers with poor economic satisfaction were more likely to be depressed. The history of pregnancy loss and pregnancy complications was another risk factor for PPD.

The aim of this study was to assess PPD risk factors. By identifying PPD risk factors in each area, preventive and curative interventions will be more effective.

Acknowledgment

The authors would like to thank the community health centers staff for cooperation in collecting the data and mothers who participated in our study.

Conflict of Interest

Authors have no conflict of interests that could inappropriately influence this work.

References

1. Patel M, Bailey RK, Jabeen S, Ali S, Barker NC, Osiezagha K. Postpartum Depression: A

Review. *J Health Care Poor Underserved* 2012; 23: 534–542.

2. Sadock BJ, Sadock VA. Kaplan & Sadock's Concise Textbook of Clinical Psychiatry. 10th eds. Lippincott Williams & Wilkins; 2008.
3. Ellsworth-Bowers ER, Corwin EJ. Nutrition and the psychoneuroimmunology of postpartum depression. *Nutr Res Rev* 2012; 25: 180–192.
4. Pearlstein T, Howard M, Salisbury A, Zlotnick C. Postpartum depression. *Am J Obstet Gynecol* 2009; 200: 357–364.
5. Field T. Postpartum depression effects on early interactions, parenting, and safety practices: a review. *Infant Behav Dev* 2010; 33: 1–6.
6. Ghaedrahmati M, Kazemi A, Kheirabadi G, Ebrahimi A, Bahrami M. Postpartum depression risk factors: A narrative review. *J Educ Health Promot.* 2017; 6: 60–60.
7. Veisani Y, Delpisheh A, Sayehmiri K, Rezaeian S. Trends of Postpartum Depression in Iran: A Systematic Review and Meta-Analysis. *Depress Res Treat* 2013; 2013: e291029.
8. Giannandrea SAM, Cerulli C, Anson E, Chaudron LH. Increased risk for postpartum psychiatric disorders among women with past pregnancy loss. *J Womens Health* 2002. 2013; 22: 760–768.
9. Hahn-Holbrook J, Cornwell-Hinrichs T, Anaya I. Economic and Health Predictors of National Postpartum Depression Prevalence: A Systematic Review, Meta-analysis, and Meta-Regression of 291 Studies from 56 Countries. *Front Psychiatry* 2017; 8: 248.
10. Braveman P, Marchi K, Egerter S, Kim S, Metzler M, Stancil T, et al. Poverty, near-poverty, and hardship around the time of pregnancy. *Matern Child Health J* 2010; 14: 20–35.
11. Abbasi N, Ghoochani OM, Ghanian M, Kitterlin M. Assessment of Households' Food Insecurity through use of a USDA Questionnaire. *Adv Plants Agric Res* 2016; 4:1–8.
12. Ivers LC, Cullen KA. Food insecurity: special considerations for women. *Am J Clin Nutr* 2011; 94: 1740S-1744S.
13. Weaver LJ, Hadley C. Moving Beyond Hunger and Nutrition: A Systematic Review of the Evidence Linking Food Insecurity and Mental Health in Developing Countries. *Ecol Food Nutr* 2009; 48: 263–284.

14. Bronte-Tinkew J, Zaslow M, Capps R, Horowitz A, McNamara M. Food Insecurity Works through Depression, Parenting, and Infant Feeding to Influence Overweight and Health in Toddlers. *J Nutr* 2007; 137: 2160–2165.
15. Laraia BA, Siega-Riz AM, Gundersen C, Dole N. Psychosocial Factors and Socioeconomic Indicators Are Associated with Household Food Insecurity among Pregnant Women. *J Nutr* 2006; 136: 177–182.
16. Hadley C, Patil CL. Food insecurity in rural Tanzania is associated with maternal anxiety and depression. *Am J Hum Biol* 2006; 18: 359–368.
17. Dewing S, Tomlinson M, le Roux IM, Chopra M, Tsai AC. Food insecurity and its association with co-occurring postnatal depression, hazardous drinking, and suicidality among women in peri-urban South Africa. *J Affect Disord* 2013; 150: 460–465.
18. Edalati-Fard F, Mirghafourvand M, Mohammad-Alizadeh-Charandabi S, Farshbaf-Khalili A. Relationship of Zinc and Magnesium Serum Levels with Postpartum Depression in Tabriz-Iran. *Glob J Health Sci* 2016; 8: 120-126.
19. Maracy MR, Iranpour S, Esmailzadeh A, Kheirabadi GHR. [Dietary Patterns During Pregnancy and the Risk of Postpartum Depression (In Persian)]. *Iran J Epidemiol* 2014 15; 10: 45–55.
20. Payab M, Motlagh AD, Eshraghian M, Rostami R, Siassi F. The association of family food security and depression in mothers having primary school children in Ray-Iran. *J Diabetes Metab Disord* 2014; 13: 65.
21. Bicke G, Nord M, Price C, Hamilton WL, Cook JT. Guide to measuring household food security, revised 2000. Alexandria, VA: USDA Food and Nutrition Service; 2000.
22. Ramesh T, Dorosty Motlagh AR, Abdollahi M. [Prevalence of household food insecurity in the City of Shiraz and its association with socio-economic and demographic factors, 2008 (In Persian)]. *Iran J Nutr Sci Food Technol* 2010; 4: 53–64.
23. Rafiei M, Nord M, Sadeghizadeh A, Entezari MH. Assessing the internal validity of a household survey-based food security measure adapted for use in Iran. *Nutr J* 2009 26; 8: 28.
24. Montazeri A, Torkan B, Omidvari S. The Edinburgh Postnatal Depression Scale (EPDS): translation and validation study of the Iranian version. *BMC Psychiatry* 2007; 7: 11.
25. Tashakori A, Behbahani AZ, Irani RD. Comparison Of Prevalence Of Postpartum Depression Symptoms Between Breastfeeding Mothers And Non-breastfeeding Mothers. *Iran J Psychiatry* 2012; 7: 61–65.
26. G. Khooshemehry G, Shariati Feizabady A, Naserkhaki V. [Prevalence of Postpartum Depression and the Factors that Decides in Clinics in the North of Tehran (In Persian)]. *Sci J Hamadan Nurs Midwifery Fac* 2011; 19: 59–69.
27. Manshoori A, Bidaki R, Esmaily H. Frequency of postpartum depression and its related factors in women referred to health centers in Rafsanjan, Iran, in 2015. *J Occup Health Epidemiol* 2017; 6: 17–24.
28. Abdollahi F, Agajani-Delavar M, Zarghami M, Lye M-S. Postpartum Mental Health in First-Time Mothers: A Cohort Study. *Iran J Psychiatry Behav Sci* 2016; 10: e426.
29. Veisani Y, Sayehmiri K. Prevalence of Postpartum Depression in Iran - A Systematic Review and Meta-Analysis. *Koomesh* 2012; 15: 21–29.
30. Gausia K, Ryder D, Ali M, Fisher C, Moran A, Koblinsky M. Obstetric complications and psychological well-being: experiences of Bangladeshi women during pregnancy and childbirth. *J Health Popul Nutr* 2012; 30: 172–180.
31. Blom EA, Jansen PW, Verhulst FC, Hofman A, Raat H, Jaddoe VWV, et al. Perinatal complications increase the risk of postpartum depression. *The Generation R Study. BJOG* 2010; 117: 1390–1398.
32. Shivalli S, Gururaj N. Postnatal depression among rural women in South India: do socio-demographic, obstetric and pregnancy outcome have a role to play? *PloS One* 2015; 10: e0122079.
33. Sooki Z, Tagharobi Z, Sharifi S, Sharifi K, Musavi S, Zahmatkesh B. Predictors of Postpartum Depression in an Iranian Population, 2006. *Nurs Midwifery Stud* 2012; 1: 12–15.
34. Barton K, Redshaw M, Quigley MA, Carson C. Unplanned pregnancy and subsequent psychological distress in partnered women: a cross-sectional study of the role of relationship quality and wider social support. *BMC Pregnancy Childbirth*. 2017 Jan 26; 17(1):44.
35. Ria MB, Budihastuti UR, Sudiyanto A. Risk Factors of Postpartum Depression at Dr. Moewardi Hospital, Surakarta. *J Matern Child Health* 2018; 3: 81–90.
36. Shivani A, Patel PJS. Unwanted childbearing and household food insecurity in the United States. *Matern Child Nutr* 2016; 12: 362–372.
37. Patel V, Rodrigues M, DeSouza N. Gender, Poverty, and Postnatal Depression: A Study of Mothers in Goa, India. *Am J Psychiatry* 2002; 159: 43–47.
38. Hromi-Fiedler A, Bermúdez-Millán A, Segura-Pérez S, Damio G, Pérez-Escamilla R. Adaptation of the US Household Food Security Survey Module for Low-Income Pregnant Latinas: Qualitative Phase. *J Hunger Environ Nutr* 2009; 4: 62–80.
39. Qobadi M, Collier C, Zhang L. The Effect of Stressful Life Events on Postpartum Depression: Findings from the 2009–2011 Mississippi Pregnancy Risk Assessment Monitoring System. *Matern Child Health J* 2016; 20: 164–172 .

40. Gundersen C, Weinreb L, Wehler C, Hosmer D. Homelessness and food insecurity. *J Hous Econ* 2003; 12: 250–272.